

# Measurement of $t\bar{t}$ quark pair cross-section with tau lepton in final state and lepton universality test

Oleksii Toldaiev

supervised by Michele Galinaro and Joao Varela

LIP, CMS

29 June 2018






TÉCNICO  
LISBOA

**FCT**

Fundação para a Ciência e a Tecnologia  
MINISTÉRIO DA CIÊNCIA, TECNOLOGIA E ENSINO SUPERIOR




# CMS detector, available data



The LHC schedule and luminosity collected by CMS, promising perspective of  $100\text{fb}^{-1}$  of 13 TeV data at end of Run 2.


# Motivation



Cross-section measurements in  $t\bar{t}$  channels at 8 TeV CMS data from 2012.

- Measurement in  $t\bar{t} \rightarrow b\bar{b}l\tau$  channel
- Improved uncertainty in estimation of main background
- It serves as preliminary work for further measurements in similar final states

# Features of $t\bar{t} \rightarrow b\bar{b}\ell\tau$ channel, measurement method



Many particular final products:


- 2 b-jets (displaced vertex of jet)
- 1 lepton (muon or electron)
- 2 neutrinos
- 1 hadronically decaying tau lepton

- Sample of  $t\bar{t}$  events is selected with simple cuts and identification requirements.
- Main background from fake taus in  $t\bar{t} \rightarrow \ell\nu_\ell q\bar{q}$  channel.
- The events are separated into background-rich and signal-rich categories according to kinematics of jets.
- The shape fit of  $M_T(\ell, E_T^{miss})$  distributions is performed.
- Both methods constrain background of misidentified taus and cross-check each other.


# Reconstruction algorithms and event selection

Standard algorithms are employed: Particle Flow for basic objects, anti-Kt jet clustering, MVA-based b-tagging, quality requirements for muons and electrons, MVA-based tau ID etc.

Require: 1 lepton,  $\geq 3$  jets,  $\geq 1$  b-tagged and 1 tau lepton.




no tau requirement




tau of Opposite Sign to muon

# Background of misidentified taus

No loss of kinematic information in  $W \rightarrow q\bar{q}$  of the background  $\ell j$  provides separation between this background and signal via the shape of transverse mass distributions  $M_T(\ell, E_T^{miss})$  and kinematic difference in jets.



transverse mass  
 $M_T(\ell, E_T^{miss})$



masses of jet combinations for W and t mass  
constraint in signal (left) and background (right)

# Profile Likelihood Ratio (PLR) shape fit in two categories

- Background- or signal-rich categories are defined by jet kinematic parameter. Profile likelihood ratio fit is performed in bins of  $M_T$  distribution.
- Likelihood function includes per-bin yields and systematic uncertainties as constraint nuisance parameters:


$$\mathcal{L}(\boldsymbol{\mu}, \theta_i) = \prod_k \mathcal{P}_{oisson} \left[ N_k | \hat{N}_k(\boldsymbol{\mu}, \theta_i) \right] \cdot \prod_i pdf(\theta_i, 0, 1) \quad (1)$$

- Based on the likelihood function the PLR test statistic is defined:


$$\lambda(\boldsymbol{\mu}) = \frac{\mathcal{L}(\boldsymbol{\mu}, \hat{\hat{\theta}}_i(\boldsymbol{\mu}))}{\mathcal{L}(\hat{\boldsymbol{\mu}}, \hat{\theta}_i)} \quad (2)$$

— scans over  $\lambda(\boldsymbol{\mu})$  provide estimation of uncertainties.

# Example of fit in both $e\tau_h$ and $\mu\tau_h$ in MC



Scan of signal strength.



Impacts of uncertainties on signal strength.

Results show agreement with SM and uncertainty of about 6-7% in both channels. Plots show simultaneous fit over both channels. The largest uncertainty is 5% from Tau ID.



# Lepton universality in W decays

$\Gamma(\tau^+\nu)/\Gamma(e^+\nu)$					$\Gamma_4/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>1.046 \pm 0.023</math> OUR FIT</b>					
$0.961 \pm 0.061$	980	<sup>42</sup> ABBOTT	00D D0	$E_{cm}^{pp} = 1.8$ TeV	
$0.94 \pm 0.14$	179	<sup>43</sup> ABE	92E CDF	$E_{cm}^{pp} = 1.8$ TeV	
$1.04 \pm 0.08 \pm 0.08$	754	<sup>44</sup> ALITTI	92F UA2	$E_{cm}^{pp} = 630$ GeV	
$1.02 \pm 0.20 \pm 0.12$	32	ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.995 \pm 0.112 \pm 0.083$	198	ALITTI	91C UA2	Repl. by ALITTI 92F	
$1.02 \pm 0.20 \pm 0.10$	32	ALBAJAR	87 UA1	Repl. by ALBAJAR 89	


$W \rightarrow e\nu$	$(10.75 \pm 0.13) \%$
$W \rightarrow \mu\nu$	$(10.57 \pm 0.15) \%$
$W \rightarrow \tau\nu$	$(11.25 \pm 0.20) \%$

from Particle Data Group (2012)

- excess of about  $2.5\sigma$  in current measurements:
  - performed at LEP in WW channel (off mass shell)
  - Tevatron in W+jets (triggering on tau)
  - relative uncertainty  $\approx 3.5\%$
- at LHC:
  - enough energy for on-shell dibosons and  $t\bar{t}$
  - similar measurements (like searches for charged Higgs) lack precision (about 6-10%, when 2% needed)
  - with the amount of luminosity we can sacrifice efficiency for purity

# Lepton universality test in $t\bar{t}$

The goal is to measure precisely the ratio  $\frac{W \rightarrow \tau \nu}{W \rightarrow \ell \nu}$  within  $t\bar{t}$  decay:



Roughly the ratio of the two channels looks like:

$$\begin{aligned}\sigma(\mu\tau_h) &= \sigma_{pp}(t\bar{t})B(W \rightarrow \mu)B(W \rightarrow \tau \rightarrow \tau_h) \\ \sigma(\mu\mu) &= \sigma_{pp}(t\bar{t})B(W \rightarrow \mu)(B(W \rightarrow \mu) + B(W \rightarrow \tau \rightarrow \mu))\end{aligned}\quad (3)$$


$$\frac{\sigma(\mu\tau_h)}{\sigma(\mu\mu)} = \frac{B(W \rightarrow \tau \rightarrow \tau_h)}{B(W \rightarrow \mu) + B(W \rightarrow \tau \rightarrow \mu)} = \frac{\frac{B(W \rightarrow \tau)}{B(W \rightarrow \mu)}B(\tau \rightarrow \tau_h)}{1 + \frac{B(W \rightarrow \tau)}{B(W \rightarrow \mu)}B(\tau \rightarrow \mu)}\quad (4)$$

The ratio cancels most of systematic uncertainties. But the remaining uncertainty due to tau ID is big (about 5%).

# Approaches to the measurement

Fit many channels in one model:

- constraint from simultaneous fit with different number of taus (both hadronic and leptonic) in final state
- consider kinematic difference between  $\ell$  and  $\tau_\ell$



Get pure sample of hadronic taus and add DY in  $\frac{\sigma(t\bar{t} \rightarrow \mu\tau_h)}{\sigma(t\bar{t} \rightarrow \mu\mu)} / \frac{\sigma(DY \rightarrow \mu\mu)}{\sigma(DY \rightarrow \tau_\mu\tau_h)}$ :

- aggressively cut on tau ID
- consider tau fakes in Opposite Sign and Same Sign event selections
- use  $3\pi$  tau decays and cut on high significance Secondary Vertex